

Fanfare™ Trailing Impatiens

Propagation

- Choose a well-drained medium with an EC of 0.75 to 0.80 mmhos and a pH of 5.6 to 5.9.
- Stick cuttings within 12 to 24 hours of arrival. Cuttings can be stored overnight, if necessary, at 45 to 50°F (7 to 10°C).
- Soil temperature should be maintained at 68 to 74°F (20 to 23°C) until roots are visible.
- As soon as is practical, mist should be reduced and then removed from **Fanfare** Trailing Impatiens. This will help decrease stretch of the rooted cutting.
- Begin fertilization with 75 to 100 ppm N when roots become visible. Increase to 150 ppm N as roots develop.
- **Fanfare** Trailing Impatiens will not require pinching during propagation.
- As the rooted cuttings develop, high light, appropriate water stress and moderate air temperatures should reduce the need for chemical plant growth regulators (PGR).
- Under low light and warm temperature conditions, cuttings of **Fanfare** Trailing Impatiens will stretch while in propagation. Growers may need to apply PGRs during propagation to further control growth. Bonzi applied as a 2 to 5 ppm heavy spray 4 to 6 days after sticking will decrease stem stretch.
- **Fanfare** Trailing Impatiens rooted cuttings should be ready for transplanting 21 to 24 days after sticking.

Growing On to Finish

Media

Use a soilless medium with good aeration, drainage and water-holding capacity, and a pH of 5.8 to 6.2.

Temperature

- Nights: 60 to 65°F (16 to 18°C)
- Days: 68 to 80°F (20 to 27°C)
- Night temperatures above 70°F (21°C) will delay flowering dramatically and reduce plant quality.

Light

- **Fanfare** Trailing Impatiens are daylength-neutral and will flower year-round.
- Plants grow best under moderate light intensity. The ideal range would be 4,000 to 6,000 f.c. (40,000 to 60,000 Lux).
- Plants will stretch at light intensities below 3,000 f.c. (30,000 Lux).
- Reduce light intensity when temperatures are high to prevent flower and leaf burning.

- HID lighting can be used to reduce crop time in areas with naturally low light levels.

Watering

- Allow plants to dry slightly between waterings in the first half of the crop cycle. Do not allow the plants to wilt during this period, as finished quality will be reduced.
- As the crop matures and begins to bud up and flower, avoid water stress entirely and remember that the crop will need more frequent irrigation.
- Excessive water stress will cause leaf edge damage as well as bud and flower drop.

Fertilizer

- Maintain constant fertilization at 200 ppm N.
- Excessive phosphorous and ammoniacal nitrogen will promote unwanted vegetative growth. Both should be provided in very limited quantities.
- Leach pots periodically with clear water to avoid build-up of salts.
- Controlled-release fertilizer can be used to supplement a liquid feed program.

Pinching

- **Fanfare** Trailing Impatiens are naturally self-branching and *do not require pinching*.
- Pinching will delay flowering approximately 10 to 14 days.

Controlling Growth

- Grow plants with adequate light and space.
- Avoid high ammonium and phosphorus fertilizers, and do not over-water.
- **Fanfare** Trailing Impatiens will generally flower and be saleable well before any plant growth regulators are needed.
- Bonzi (5 to 15 ppm) applied as a spray 1 to 2 times can be used to control growth of **Fanfare** Trailing Impatiens.
- **Fanfare Orange** may require more frequent and earlier application of plant growth regulator due to its more vigorous habit.
- Mature plants which are approaching shipping can be drenched with Bonzi (0.25 to 0.50 ppm) to significantly slow vegetative growth while allowing flowering to continue.
- These recommendations for plant growth regulators should be used only as general guidelines. Growers must trial all chemicals under their particular conditions.

Common Problems

Insects: Thrips, spider mites, aphids.

Diseases: Impatiens Necrotic Spot Virus (INSV), *Botrytis* (gray mold).

All **Fanfare** Trailing Impatiens cuttings are derived from culture and virus-indexed stock from the **Ball Certified Plants**[®] program.

Problems	Causes
Plants collapse	Stem canker (<i>Botrytis</i>) Plants grown in saturated soil for extended period of time (<i>Pythium</i>)
Excessive vegetative growth, lack of flowers	Excessive nitrogen in fertilizer Excessive phosphorous Over-fertilization under low light conditions Low light and over-watering; wet media Excess or late Florel application
Foliage necrosis, leaf spot	Drying out between waterings Excess minor nutrient levels in media
Poor branching, thin plants	Low fertilization in early stages of crop

Fanfare Trailing Impatiens Crop Schedule & Uses

(Crop Schedule In Weeks)

	4-In. (10-Cm) Pots 1 PP*	6-In. (15-Cm) Pots 1 PP*	10-12-In. (25-30-Cm) Pots 3-5 PP*
Unrooted cuttings	9-11	11-13	14-15
Rooted cuttings	6-8	8-10	11-13

*PP: Plants per pot or basket

Note: Growers should use the information presented here as a starting point. Crop times will vary depending on the climate, location, time of year and greenhouse environmental conditions.

Visit **Ball FloraPlant.com** today to check out the complete line of terrific products from Ball FloraPlant.

 Ball FloraPlant

In North America, Ball FloraPlant products are distributed exclusively by Ball Seed Co., Ball Superior and ColorLink.
800 879-BALL Fax: 800 234-0370 ballseed.com

[™] denotes a trademark of and [®] denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.

© 2006 Ball Horticultural Company Printed In USA BFP06052 12/05